

NAMES OF MEDICATION & PURPOSE

ORAL MEDICATIONS

___	DOXYCYCLINE	Antibiotic used to prevent infection from a procedure or minor surgery. Unless otherwise prescribed 1 Tab twice daily.
___	BABY ASPIRIN	To increase blood supply to the uterus and or ovary. Help prevent blood clotting. 1 Tab per day
___	ESTRACE	Used to thicken the lining of the uterus for transfer of embryos. 1 or more Tabs per day
___	PREDNISON	Steroid – Suppresses male hormones produced by adrenals and also suppresses immune function, which at times is needed. (10-60-10) You will start on 10 mg a day. The day after you are told to do HCG injection you are to increase dose to 60 mg for 4 days, then decrease to 1 tab daily. (10mg) again until told otherwise
___	PROVERA	Used to induce menses from 2 days-2 weeks. 1 tab two times a day for 5 days
___	SEROPHENE	(CLOMID) Enhances quality and quantity of ovulation to ensure proper timing of intercourse or artificial insemination. Used on Days 5-9 of cycle in dosage of 1-3 Tabs (50-150 mg) daily

PROGESTRONE MEDICATIONS

___	CRINONE GEL	Used to thicken the lining of the uterus. Vaginally inserted. Amount varies.
___	PROMETRIUM	Used to thicken the lining of the uterus. Vaginally inserted 2 tabs three times a day – rest 10-15 min. after insertion.
___	PROGESTRONE IN OIL	Used to thicken the lining of the uterus. Intramuscular injection. Amount varies.

SUBCUTANEOUS INJECTIONS (S.C.)

___	ANTAGON	Prevents early ovulation (Starts on Day 6 of stimulation if follicles are 12-13mm). Is given S.C.
___	LUPRON	Suppresses your hormones (body is in menopausal state) and prevents early ovulation. Usually starts day 2/3 or 21/22 of menstrual cycle. Is given S.C.
___	HEPARIN	Used to prevent recurrent pregnancy losses with antiphospholipid antibodies. (Blood thinner) also used in hyperstimulation syndrome. Is given S.C., usually at a dosage of 5000 U twice daily.
___	FOLLISTIM	Used for stimulation of follicular growth and ovulation. Given S.C. in varying dosage.
___	GONAL-F	Used for stimulation of follicular growth and ovulation. Given S.C. in varying dosage.

INTRAMUSCULAR INJECTIONS (I.M.)

___	REPRONEX	Used for stimulation of follicular growth and ovulation. Usually given I.M. in varying dosages. Can, however, also be given S.C.
___	PERGONAL	Used for stimulation of follicular growth and ovulation. Usually given I.M. in varying dosages.
___	PROFASI	(HCG) Triggers ovulation within 36 hours after injection. Given I.M. in dosage of 5000 or 10,000 IV.
___	PREGNYL	(HCG) Triggers ovulation within 36 hours after injection. Given I.M. in dosages of 5000 or 10,000 IV.